



Data From Phase 2 PALLET Study of IBRANCE® (palbociclib) in Early Breast Cancer to be Presented at San Antonio Breast Cancer Symposium

Friday, November 09, 2018 - 02:57pm

- - -

Abstract Now Available Online

NEW YORK, N.Y.– Pfizer Inc. (NYSE:PFE) today announced that results from the Phase 2 PALLET (NCT02296801) study of IBRANCE® (palbociclib) will be presented at the 41st Annual San Antonio Breast Cancer Symposium (SABCS). The abstract [#GS3-02] is now accessible on the SABCS website here ahead of the oral presentation on December 6.

PALLET is a Phase 2 randomized study evaluating the biological and clinical effects of the combination of IBRANCE with letrozole as neoadjuvant therapy in post-menopausal women with estrogen-receptor positive primary breast cancer. The study is sponsored and being conducted by the National Surgical Adjuvant Breast and Bowel Project (NSABP) Foundation Inc. in North America and The Institute of Cancer Research, London, and The Royal Marsden NHS Foundation Trust in the United Kingdom. Pfizer has provided research funding and drug supply.

About IBRANCE® (palbociclib) 125 mg capsules IBRANCE is an oral inhibitor of CDKs 4 and 6,¹ which are key regulators of the cell cycle that trigger cellular progression.^{2,3} In the U.S., IBRANCE is indicated for the treatment of HR+, HER2- advanced or metastatic breast cancer in combination with an aromatase inhibitor as initial endocrine based

therapy in postmenopausal women, or fulvestrant in women with disease progression following endocrine therapy.

The full prescribing information for IBRANCE can be found at www.pfizer.com.

IMPORTANT IBRANCE® (palbociclib) SAFETY INFORMATION FROM THE U.S. PRESCRIBING INFORMATION

Neutropenia was the most frequently reported adverse reaction in PALOMA-2 (80%) and PALOMA-3 (83%). In PALOMA-2, Grade 3 (56%) or 4 (10%) decreased neutrophil counts were reported in patients receiving IBRANCE plus letrozole. In PALOMA-3, Grade 3 (55%) or Grade 4 (11%) decreased neutrophil counts were reported in patients receiving IBRANCE plus fulvestrant. Febrile neutropenia has been reported in 1.8% of patients exposed to IBRANCE across PALOMA-2 and PALOMA-3. One death due to neutropenic sepsis was observed in PALOMA-3. Inform patients to promptly report any fever.

Monitor complete blood count prior to starting IBRANCE, at the beginning of each cycle, on Day 15 of first 2 cycles and as clinically indicated. Dose interruption, dose reduction, or delay in starting treatment cycles is recommended for patients who develop Grade 3 or 4 neutropenia.

Based on the mechanism of action, IBRANCE can cause **fetal harm**. Advise females of reproductive potential to use effective contraception during IBRANCE treatment and for at least 3 weeks after the last dose. IBRANCE may **impair fertility in males** and has the potential to cause genotoxicity. Advise male patients with female partners of reproductive potential to use effective contraception during IBRANCE treatment and for 3 months after the last dose. Advise females to inform their healthcare provider of a known or suspected pregnancy. Advise women **not to breastfeed** during IBRANCE treatment and for 3 weeks after the last dose because of the potential for serious adverse reactions in nursing infants.

The **most common adverse reactions (≥10%)** of any grade reported in **PALOMA-2** for IBRANCE plus letrozole vs placebo plus letrozole were neutropenia (80% vs 6%), infections (60% vs 42%), leukopenia (39% vs 2%), fatigue (37% vs 28%), nausea (35% vs 26%), alopecia (33% vs 16%), stomatitis (30% vs 14%), diarrhea (26% vs 19%), anemia (24% vs 9%), rash (18% vs 12%), asthenia (17% vs 12%), thrombocytopenia (16% vs 1%), vomiting (16% vs 17%), decreased appetite (15% vs 9%), dry skin (12% vs 6%), pyrexia (12% vs 9%), and dysgeusia (10% vs 5%).

The **most frequently reported Grade ≥ 3 adverse reactions ($\geq 5\%$)** in **PALOMA-2** for IBRANCE plus letrozole vs placebo plus letrozole were neutropenia (66% vs 2%), leukopenia (25% vs 0%), infections (7% vs 3%), and anemia (5% vs 2%). Lab abnormalities of any grade occurring in PALOMA-2 for IBRANCE plus letrozole vs placebo plus letrozole were decreased WBC (97% vs 25%), decreased neutrophils (95% vs 20%), anemia (78% vs 42%), decreased platelets (63% vs 14%), increased aspartate aminotransferase (52% vs 34%), and increased alanine aminotransferase (43% vs 30%).

The **most common adverse reactions ($\geq 10\%$)** of any grade reported in **PALOMA-3** for IBRANCE plus fulvestrant vs placebo plus fulvestrant were neutropenia (83% vs 4%), leukopenia (53% vs 5%), infections (47% vs 31%), fatigue (41% vs 29%), nausea (34% vs 28%), anemia (30% vs 13%), stomatitis (28% vs 13%), diarrhea (24% vs 19%), thrombocytopenia (23% vs 0%), vomiting (19% vs 15%), alopecia (18% vs 6%), rash (17% vs 6%), decreased appetite (16% vs 8%), and pyrexia (13% vs 5%).

The **most frequently reported Grade ≥ 3 adverse reactions ($\geq 5\%$)** in **PALOMA-3** for IBRANCE plus fulvestrant vs placebo plus fulvestrant were neutropenia (66% vs 1%) and leukopenia (31% vs 2%).

Lab abnormalities of any grade occurring in **PALOMA-3** for IBRANCE plus fulvestrant vs placebo plus fulvestrant were decreased WBC (99% vs 26%), decreased neutrophils (96% vs 14%), anemia (78% vs 40%), decreased platelets (62% vs 10%), increased aspartate aminotransferase (43% vs 48%), and increased alanine aminotransferase (36% vs 34%).

Avoid concurrent use of **strong CYP3A inhibitors**. If patients must be administered a strong CYP3A inhibitor, reduce the IBRANCE dose to 75 mg. If the strong inhibitor is discontinued, increase the IBRANCE dose (after 3-5 half-lives of the inhibitor) to the dose used prior to the initiation of the strong CYP3A inhibitor. Grapefruit or grapefruit juice may increase plasma concentrations of IBRANCE and should be avoided. Avoid concomitant use of **strong CYP3A inducers**. The dose of **sensitive CYP3A substrates** with a narrow therapeutic index may need to be reduced as IBRANCE may increase their exposure.

For patients with **severe hepatic impairment** (Child-Pugh class C), the recommended dose of IBRANCE is 75 mg. The pharmacokinetics of IBRANCE **have not been studied** in patients **requiring hemodialysis**.

About Pfizer Oncology At Pfizer Oncology, we are committed to advancing medicines wherever we believe we can make a meaningful difference on the lives of patients.

Today, Pfizer Oncology has an industry-leading portfolio of 13 approved cancer medicines across 21 indications, including breast, prostate, kidney, lung and hematology. We also have one of the deepest oncology biosimilars pipelines, with two medicines approved globally and several assets in mid to late-stage development for the treatment of cancer or as supportive care. Pfizer Oncology is striving to change the trajectory of cancer.

Working together for a healthier world® At Pfizer, we apply science and our global resources to bring therapies to people that extend and significantly improve their lives. We strive to set the standard for quality, safety and value in the discovery, development and manufacture of health care products. Our global portfolio includes medicines and vaccines as well as many of the world's best-known consumer health care products. Every day, Pfizer colleagues work across developed and emerging markets to advance wellness, prevention, treatments and cures that challenge the most feared diseases of our time. Consistent with our responsibility as one of the world's premier innovative biopharmaceutical companies, we collaborate with health care providers, governments and local communities to support and expand access to reliable, affordable health care around the world. For more than 150 years, we have worked to make a difference for all who rely on us. We routinely post information that may be important to investors on our website at www.pfizer.com. In addition, to learn more, please visit us on www.pfizer.com and follow us on Twitter at @Pfizer and @Pfizer_News, LinkedIn, YouTube, and like us on Facebook at [Facebook.com/Pfizer](https://www.facebook.com/Pfizer).

DISCLOSURE NOTICE: The information contained in this release is as of November 9, 2018. Pfizer assumes no obligation to update forward-looking statements contained in this release as the result of new information or future events or developments. This release contains forward-looking information about IBRANCE (palbociclib), including its potential benefits, that involves substantial risks and uncertainties that could cause actual results to differ materially from those expressed or implied by such statements. Risks and uncertainties include, among other things, uncertainties regarding the commercial success of IBRANCE; the uncertainties inherent in research and development, including the ability to meet anticipated clinical trial commencement and completion dates and regulatory submission dates, as well as the possibility of unfavorable clinical trial results, including unfavorable new clinical data and additional analyses of existing clinical data; whether regulatory authorities will be satisfied with the design of and results from our clinical studies; whether and when drug applications may be filed in any jurisdictions for IBRANCE for potential early breast cancer indications or in any jurisdictions for any other potential indications for IBRANCE; whether and when any such applications may be approved by regulatory authorities, which will depend on the assessment by such regulatory authorities of the benefit-risk profile suggested by the

totality of the efficacy and safety information submitted; decisions by regulatory authorities regarding labeling and other matters that could affect the availability or commercial potential of IBRANCE; and competitive developments. A further description of risks and uncertainties can be found in Pfizer's Annual Report on Form 10-K for the fiscal year ended December 31, 2017 and in its subsequent reports on Form 10-Q, including in the sections thereof captioned "Risk Factors" and "Forward-Looking Information and Factors That May Affect Future Results," as well as in its subsequent reports on Form 8-K, all of which are filed with the U.S. Securities and Exchange Commission and available at www.sec.gov and www.pfizer.com. # # # # # 1

IBRANCE® (palbociclib) Prescribing Information. New York. NY: Pfizer Inc: 2018.2

Weinberg RA. pRb and Control of the Cell Cycle Clock. In: Weinberg RA, ed. The Biology of Cancer. 2nd ed. New York, NY: Garland Science; 2014:275- 329.3

Sotillo E, Grana X. Escape from Cellular Quiescence. In: Enders GH, ed. Cell Cycle Deregulation in Cancer. New York, NY: Humana Press; 2010:3-22. Pfizer U.S. Media Contact: Jessica Smith (212)733-6213 Jessica.M.Smith@pfizer.com Pfizer Investor Contact: Ryan Crowe (212)733-8160 Ryan.Crowe@pfizer.com