



# European Medicines Agency Validates Application for BAVENCIO® (avelumab) Plus INLYTA® (axitinib) for the Treatment of Advanced Renal Cell Carcinoma

Friday, March 08, 2019 - 02:00am

Merck KGaA, Darmstadt, Germany, which operates its biopharmaceutical business as EMD Serono in the US and Canada, and Pfizer Inc. (NYSE: PFE) today announced that the European Medicines Agency (EMA) has validated for review the Type II variation application for BAVENCIO® (avelumab) in combination with INLYTA® (axitinib)\* for the treatment of patients with advanced renal cell carcinoma (RCC). With this validation, the application is complete, and the EMA will now begin the review procedure.

The application is based on results from the pivotal Phase III JAVELIN Renal 101 trial, which were published in the *New England Journal of Medicine* on February 16, 2019.

The US Food and Drug Administration has also accepted a supplemental Biologics License Application for BAVENCIO in combination with INLYTA for patients with advanced RCC for Priority Review, with a target action date in June 2019. A supplemental application for the combination in unresectable or metastatic RCC has also been submitted in Japan.

Despite available therapies, the outlook for patients with advanced RCC remains poor.<sup>1</sup> Approximately 20% to 30% of patients are first diagnosed at the metastatic stage.<sup>2</sup> The five-year survival rate for patients with metastatic RCC is approximately 12%.<sup>1</sup>

The clinical development program for avelumab, known as JAVELIN, involves at least 30 clinical programs and more than 9,000 patients evaluated across more than 15 different

tumor types. In addition to RCC, these tumor types include breast, gastric/gastro-esophageal junction, and head and neck cancers, Merkel cell carcinoma, non-small cell lung cancer, and urothelial carcinoma.

\*The combination of BAVENCIO and INLYTA is under clinical investigation for advanced RCC, and there is no guarantee this combination will be approved for advanced RCC by any health authority worldwide. INLYTA is approved by the EMA for use in the EU in adult patients with advanced RCC after failure of prior treatment with sunitinib or a cytokine. In the US, INLYTA is approved as monotherapy for the treatment of advanced RCC after failure of one prior systemic therapy.

**About Renal Cell Carcinoma** RCC is the most common form of kidney cancer, accounting for about 2% to 3% of all cancers in adults.<sup>3,4</sup> The most common type of RCC is clear cell carcinoma, accounting for approximately 70% of all cases.<sup>3</sup> About 115,174 kidney, renal pelvis and ureter cancers were diagnosed in Europe in 2012, accounting for 3.9% of all new cancer cases in Europe.<sup>5</sup> An estimated 73,820 new cases of kidney cancer will be diagnosed in the US in 2019, and approximately 14,770 people will die from the disease.<sup>6</sup>

**About BAVENCIO® (avelumab)** BAVENCIO is a human anti-programmed death ligand-1 (PD-L1) antibody. BAVENCIO has been shown in preclinical models to engage both the adaptive and innate immune functions. By blocking the interaction of PD-L1 with PD-1 receptors, BAVENCIO has been shown to release the suppression of the T cell-mediated antitumor immune response in preclinical models.<sup>7-9</sup> BAVENCIO has also been shown to induce NK cell-mediated direct tumor cell lysis via antibody-dependent cell-mediated cytotoxicity (ADCC) in vitro.<sup>9-11</sup> In November 2014, Merck KGaA, Darmstadt, Germany, and Pfizer announced a strategic alliance to co-develop and co-commercialize BAVENCIO.

### **Approved Indications in the US**

In the US, the FDA granted accelerated approval for BAVENCIO for the treatment of (i) adults and pediatric patients 12 years and older with metastatic Merkel cell carcinoma (mMCC) and (ii) patients with locally advanced or metastatic urothelial carcinoma (mUC) who have disease progression during or following platinum-containing chemotherapy, or have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy. These indications are approved under accelerated approval based on tumor response rate and duration of response. Continued approval for these indications may be contingent upon verification and description of clinical benefit in confirmatory trials.

BAVENCIO is currently approved for patients with MCC in more than 45 countries globally, with the majority of these approvals in a broad indication that is not limited to a specific line of treatment.

### **Important Safety Information from the US FDA-Approved Label**

BAVENCIO can cause **immune-mediated pneumonitis**, including fatal cases. Monitor patients for signs and symptoms of pneumonitis, and evaluate suspected cases with radiographic imaging. Administer corticosteroids for Grade 2 or greater pneumonitis. Withhold BAVENCIO for moderate (Grade 2) and permanently discontinue for severe (Grade 3), life-threatening (Grade 4), or recurrent moderate (Grade 2) pneumonitis. Pneumonitis occurred in 1.2% (21/1738) of patients, including one (0.1%) patient with Grade 5, one (0.1%) with Grade 4, and five (0.3%) with Grade 3.

BAVENCIO can cause **immune-mediated hepatitis**, including fatal cases. Monitor patients for abnormal liver tests prior to and periodically during treatment. Administer corticosteroids for Grade 2 or greater hepatitis. Withhold BAVENCIO for moderate (Grade 2) immune-mediated hepatitis until resolution and permanently discontinue for severe (Grade 3) or life-threatening (Grade 4) immune-mediated hepatitis. Immune-mediated hepatitis was reported in 0.9% (16/1738) of patients, including two (0.1%) patients with Grade 5, and 11 (0.6%) with Grade 3.

BAVENCIO can cause **immune-mediated colitis**. Monitor patients for signs and symptoms of colitis. Administer corticosteroids for Grade 2 or greater colitis. Withhold BAVENCIO until resolution for moderate or severe (Grade 2 or 3) colitis, and permanently discontinue for life-threatening (Grade 4) or recurrent (Grade 3) colitis upon reinitiation of BAVENCIO. Immune-mediated colitis occurred in 1.5% (26/1738) of patients, including seven (0.4%) with Grade 3.

BAVENCIO can cause **immune-mediated endocrinopathies**, including adrenal insufficiency, thyroid disorders, and type 1 diabetes mellitus.

Monitor patients for signs and symptoms of **adrenal insufficiency** during and after treatment, and administer corticosteroids as appropriate. Withhold BAVENCIO for severe (Grade 3) or life-threatening (Grade 4) adrenal insufficiency. Adrenal insufficiency was reported in 0.5% (8/1738) of patients, including one (0.1%) with Grade 3.

**Thyroid disorders** can occur at any time during treatment. Monitor patients for changes in thyroid function at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation. Manage hypothyroidism with hormone

replacement therapy and hyperthyroidism with medical management. Withhold BAVENCIO for severe (Grade 3) or life-threatening (Grade 4) thyroid disorders. Thyroid disorders, including hypothyroidism, hyperthyroidism, and thyroiditis, were reported in 6% (98/1738) of patients, including three (0.2%) with Grade 3.

**Type 1 diabetes mellitus** including diabetic ketoacidosis: Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Withhold BAVENCIO and administer antihyperglycemics or insulin in patients with severe or life-threatening (Grade  $\geq$  3) hyperglycemia, and resume treatment when metabolic control is achieved. Type 1 diabetes mellitus without an alternative etiology occurred in 0.1% (2/1738) of patients, including two cases of Grade 3 hyperglycemia.

BAVENCIO can cause **immune-mediated nephritis and renal dysfunction**. Monitor patients for elevated serum creatinine prior to and periodically during treatment. Administer corticosteroids for Grade 2 or greater nephritis. Withhold BAVENCIO for moderate (Grade 2) or severe (Grade 3) nephritis until resolution to Grade 1 or lower. Permanently discontinue

BAVENCIO for life-threatening (Grade 4) nephritis. Immune-mediated nephritis occurred in 0.1% (1/1738) of patients.

BAVENCIO can result in **other severe and fatal immune-mediated adverse reactions** involving any organ system during treatment or after treatment discontinuation. For suspected immune-mediated adverse reactions, evaluate to confirm or rule out an immune-mediated adverse reaction and to exclude other causes. Depending on the severity of the adverse reaction, withhold or permanently discontinue BAVENCIO, administer high-dose corticosteroids, and initiate hormone replacement therapy, if appropriate. Resume BAVENCIO when the immune-mediated adverse reaction remains at Grade 1 or lower following a corticosteroid taper. Permanently discontinue BAVENCIO for any severe (Grade 3) immune-mediated adverse reaction that recurs and for any life-threatening (Grade 4) immune-mediated adverse reaction. The following clinically significant immune-mediated adverse reactions occurred in less than 1% of 1738 patients treated with BAVENCIO: myocarditis with fatal cases, myositis, psoriasis, arthritis, exfoliative dermatitis, erythema multiforme, pemphigoid, hypopituitarism, uveitis, Guillain-Barré syndrome, and systemic inflammatory response.

BAVENCIO can cause severe (Grade 3) or life-threatening (Grade 4) **infusion-related reactions**. Patients should be premedicated with an antihistamine and acetaminophen prior to the first 4 infusions and for subsequent doses based upon clinical judgment and

presence/severity of prior infusion reactions. Monitor patients for signs and symptoms of infusion-related reactions, including pyrexia, chills, flushing, hypotension, dyspnea, wheezing, back pain, abdominal pain, and urticaria. Interrupt or slow the rate of infusion for mild (Grade 1) or moderate (Grade 2) infusion-related reactions. Permanently discontinue BAVENCIO for severe (Grade 3) or life-threatening (Grade 4) infusion-related reactions. Infusion-related reactions occurred in 25% (439/1738) of patients, including three (0.2%) patients with Grade 4 and nine (0.5%) with Grade 3.

BAVENCIO can cause **fetal harm** when administered to a pregnant woman. Advise patients of the potential risk to a fetus including the risk of fetal death. Advise females of childbearing potential to use effective contraception during treatment with BAVENCIO and for at least 1 month after the last dose of BAVENCIO. It is not known whether BAVENCIO is excreted in human milk. Advise a lactating woman **not to breastfeed** during treatment and for at least 1 month after the last dose of BAVENCIO due to the potential for serious adverse reactions in breastfed infants.

**The most common adverse reactions** (all grades,  $\geq 20\%$ ) in patients with **metastatic Merkel cell carcinoma (MCC)** were fatigue (50%), musculoskeletal pain (32%), diarrhea (23%), nausea (22%), infusion-related reaction (22%), rash (22%), decreased appetite (20%), and peripheral edema (20%).

**Selected treatment-emergent laboratory abnormalities** (all grades,  $\geq 20\%$ ) in patients with **metastatic MCC** were lymphopenia (49%), anemia (35%), increased aspartate aminotransferase (34%), thrombocytopenia (27%), and increased alanine aminotransferase (20%).

**The most common adverse reactions** (all grades,  $\geq 20\%$ ) in patients with **locally advanced or metastatic urothelial carcinoma (UC)** were fatigue (41%), infusion-related reaction (30%), musculoskeletal pain (25%), nausea (24%), decreased appetite/hypophagia (21%), and urinary tract infection (21%).

**Selected laboratory abnormalities** (Grades 3-4,  $\geq 3\%$ ) in patients with **locally advanced or metastatic UC** were hyponatremia (16%), increased gamma-glutamyltransferase (12%), lymphopenia (11%), hyperglycemia (9%), increased alkaline phosphatase (7%), anemia (6%), increased lipase (6%), hyperkalemia (3%), and increased aspartate aminotransferase (3%).

Please see full US Prescribing Information and Medication Guide available at <http://www.BAVENCIO.com>.

## **About INLYTA® (axitinib)**

INLYTA is an oral therapy that is designed to inhibit tyrosine kinases, including vascular endothelial growth factor (VEGF) receptors 1, 2 and 3; these receptors can influence tumor growth, vascular angiogenesis and progression of cancer (the spread of tumors). In the US, INLYTA is approved for the treatment of advanced renal cell carcinoma (RCC) after failure of one prior systemic therapy. INLYTA is also approved by the European Medicines Agency (EMA) for use in the EU in adult patients with advanced RCC after failure of prior treatment with sunitinib or a cytokine.

**INLYTA Important Safety Information Hypertension** including **hypertensive crisis** has been observed. Blood pressure should be well controlled prior to initiating INLYTA. Monitor for hypertension and treat as needed. For persistent hypertension, despite use of antihypertensive medications, reduce the dose. Discontinue INLYTA if hypertension is severe and persistent despite use of antihypertensive therapy and dose reduction of INLYTA, and discontinuation should be considered if there is evidence of hypertensive crisis.

**Arterial and venous thrombotic events** have been observed and can be fatal. Use with caution in patients who are at increased risk or who have a history of these events.

**Hemorrhagic events**, including fatal events, have been reported. INLYTA has not been studied in patients with evidence of untreated brain metastasis or recent active gastrointestinal bleeding and should not be used in those patients. If any bleeding requires medical intervention, temporarily interrupt the INLYTA dose.

**Cardiac failure** has been observed and can be fatal. Monitor for signs or symptoms of cardiac failure throughout treatment with INLYTA. Management of cardiac failure may require permanent discontinuation of INLYTA.

**Gastrointestinal perforation and fistula**, including death, have occurred. Use with caution in patients at risk for gastrointestinal perforation or fistula. Monitor for symptoms of gastrointestinal perforation or fistula periodically throughout treatment.

**Hypothyroidism** requiring thyroid hormone replacement has been reported. Monitor thyroid function before initiation of, and periodically throughout, treatment.

No formal studies of the effect of INLYTA on **wound healing** have been conducted. Stop INLYTA at least 24 hours prior to scheduled surgery.

**Reversible Posterior Leukoencephalopathy Syndrome (RPLS)** has been observed. If signs or symptoms occur, permanently discontinue treatment.

Monitor for **proteinuria** before initiation of, and periodically throughout, treatment. For moderate to severe proteinuria, reduce the dose or temporarily interrupt treatment.

**Liver enzyme elevation** has been observed during treatment with INLYTA. Monitor ALT, AST, and bilirubin before initiation of, and periodically throughout, treatment.

For patients with moderate **hepatic impairment**, the starting dose should be decreased. INLYTA has not been studied in patients with severe hepatic impairment.

Women of childbearing potential should be advised of potential hazard to the fetus and to avoid becoming **pregnant** while receiving INLYTA.

Avoid strong **CYP3A4/5 inhibitors**. If unavoidable, reduce the dose. Grapefruit or grapefruit juice may also increase INLYTA plasma concentrations and should be avoided.

Avoid strong **CYP3A4/5 inducers** and, if possible, avoid moderate CYP3A4/5 inducers.

The **most common ( $\geq 20\%$ ) adverse events (AEs)** occurring in patients receiving INLYTA (all grades, vs sorafenib) were diarrhea (55% vs 53%), hypertension (40% vs 29%), fatigue (39% vs 32%), decreased appetite (34% vs 29%), nausea (32% vs 22%), dysphonia (31% vs 14%), hand-foot syndrome (27% vs 51%), weight decreased (25% vs 21%), vomiting (24% vs 17%), asthenia (21% vs 14%), and constipation (20% vs 20%).

The **most common ( $\geq 10\%$ ) grade 3/4 AEs** occurring in patients receiving INLYTA (vs sorafenib) were hypertension (16% vs 11%), diarrhea (11% vs 7%), and fatigue (11% vs 5%).

The **most common ( $\geq 20\%$ ) lab abnormalities** occurring in patients receiving INLYTA (all grades, vs sorafenib) included increased creatinine (55% vs 41%), decreased bicarbonate (44% vs 43%), hypocalcemia (39% vs 59%), decreased hemoglobin (35% vs 52%), decreased lymphocytes (absolute) (33% vs 36%), increased ALP (30% vs 34%), hyperglycemia (28% vs 23%), increased lipase (27% vs 46%), increased amylase (25% vs 33%), increased ALT (22% vs 22%), and increased AST (20% vs 25%).

For more information and full Prescribing Information, visit [www.INLYTA.com](http://www.INLYTA.com).

**Alliance between Merck KGaA, Darmstadt, Germany, and Pfizer Inc., New York, US** Immuno-oncology is a top priority for Merck KGaA, Darmstadt, Germany, and Pfizer.

The global strategic alliance between Merck KGaA, Darmstadt, Germany, and Pfizer enables the companies to benefit from each other's strengths and capabilities and further explore the therapeutic potential of BAVENCIO, an anti-PD-L1 antibody initially discovered and developed by Merck KGaA, Darmstadt, Germany. The immuno-oncology alliance is jointly developing and commercializing BAVENCIO. The alliance is focused on developing high-priority international clinical programs to investigate BAVENCIO as a monotherapy as well as combination regimens, and is striving to find new ways to treat cancer.

All Merck KGaA, Darmstadt, Germany, Press Releases are distributed by e-mail at the same time they become available on the Merck KGaA, Darmstadt, Germany, Website. Please go to [www.emdgroup.com/subscribe](http://www.emdgroup.com/subscribe) to register online, change your selection or discontinue this service.

**About Merck KGaA, Darmstadt, Germany** Merck KGaA, Darmstadt, Germany, a leading science and technology company, operates across healthcare, life science and performance materials. Around 51,000 employees work to make a positive difference to millions of people's lives every day by creating more joyful and sustainable ways to live. From advancing gene editing technologies and discovering unique ways to treat the most challenging diseases to enabling the intelligence of devices – the company is everywhere. In 2017, Merck KGaA, Darmstadt, Germany, generated sales of € 15.3 billion in 66 countries.

The company holds the global rights to the name and trademark "Merck" internationally. The only exceptions are the United States and Canada, where the business sectors of Merck KGaA, Darmstadt, Germany operate as EMD Serono in healthcare, MilliporeSigma in life science, and EMD Performance Materials. Since its founding in 1668, scientific exploration and responsible entrepreneurship have been key to the company's technological and scientific advances. To this day, the founding family remains the majority owner of the publicly listed company.

**Pfizer Inc.: Working together for a healthier world®** At Pfizer, we apply science and our global resources to bring therapies to people that extend and significantly improve their lives. We strive to set the standard for quality, safety and value in the discovery, development and manufacture of health care products. Our global portfolio includes medicines and vaccines as well as many of the world's best-known consumer health care products. Every day, Pfizer colleagues work across developed and emerging markets to advance wellness, prevention, treatments and cures that challenge the most feared diseases of our time. Consistent with our responsibility as one of the world's premier innovative biopharmaceutical companies, we collaborate with health care providers,



governments and local communities to support and expand access to reliable, affordable health care around the world. For more than 150 years, we have worked to make a difference for all who rely on us. We routinely post information that may be important to investors on our website at [www.pfizer.com](http://www.pfizer.com). In addition, to learn more, please visit us on [www.pfizer.com](http://www.pfizer.com) and follow us on Twitter at @Pfizer and @Pfizer\_News, LinkedIn, YouTube and like us on Facebook at [Facebook.com/Pfizer](https://www.facebook.com/Pfizer).

*Pfizer Disclosure Notice The information contained in this release is as of March 8, 2019. Pfizer assumes no obligation to update forward-looking statements contained in this release as the result of new information or future events or developments.*

*This release contains forward-looking information about BAVENCIO (avelumab), including a potential new indication for BAVENCIO in combination with INLYTA (axitinib) for the treatment of patients with advanced renal cell carcinoma, the alliance between Merck KGaA, Darmstadt, Germany, and Pfizer involving BAVENCIO and clinical development plans, including their potential benefits, that involves substantial risks and uncertainties that could cause actual results to differ materially from those expressed or implied by such statements. Risks and uncertainties include, among other things, uncertainties regarding the commercial success of BAVENCIO; the uncertainties inherent in research and development, including the ability to meet anticipated clinical endpoints, commencement and/or completion dates for our clinical trials, regulatory submission dates, regulatory approval dates and/or launch dates, as well as the possibility of unfavorable new clinical data and further analyses of existing clinical data and uncertainties regarding whether the other primary endpoint of JAVELIN Renal 101 will be met; risks associated with interim data; the risk that clinical trial data are subject to differing interpretations and assessments by regulatory authorities; whether regulatory authorities will be satisfied with the design of and results from our clinical studies; whether and when any drug applications may be filed for BAVENCIO in combination with INLYTA for the potential new indication in any other jurisdictions or in any jurisdictions for any other potential indications for BAVENCIO or combination therapies; whether and when the pending applications with the EMA, as well as in the US and Japan for BAVENCIO in combination with INLYTA for the potential new indication may be approved and whether and when regulatory authorities in any jurisdictions where any other applications are pending or may be submitted for BAVENCIO or combination therapies may approve any such applications, which will depend on myriad factors, including making a determination as to whether the product's benefits outweigh its known risks and determination of the product's efficacy, and, if approved, whether they will be commercially successful; decisions by regulatory authorities impacting labeling,*

*manufacturing processes and/or other matters that could affect the availability or commercial potential of BAVENCIO or combination therapies, including BAVENCIO in combination with INLYTA for the potential new indication; and competitive developments.*

*A further description of risks and uncertainties can be found in Pfizer's Annual Report on Form 10-K for the fiscal year ended December 31, 2018, and in its subsequent reports on Form 10-Q, including in the sections thereof captioned "Risk Factors" and "Forward-Looking Information and Factors That May Affect Future Results", as well as in its subsequent reports on Form 8-K, all of which are filed with the U.S. Securities and Exchange Commission and available at [www.sec.gov](http://www.sec.gov) and [www.pfizer.com](http://www.pfizer.com).*

## **References**

Motzer R, Penkov K, Haanen J, et al. Avelumab plus axitinib versus sunitinib for advanced renal-cell carcinoma. *NEJM*. February 16, 2019. Published online ahead of print. National Cancer Institute: SEER Stat Fact Sheets: Kidney and renal pelvis. Available from: <http://seer.cancer.gov/statfacts/html/kidrp.html>. Accessed February 2019. American Cancer Society. What is kidney cancer? Available from: <https://www.cancer.org/cancer/kidney-cancer/about.html>. Accessed February 2019. Escudier B, Porta C, Schmidinger M et al Renal cell carcinoma: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2014; 25(Suppl3):iii49-iii56. American Cancer Society. Cancer facts and figures 2019. Available at: <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2019/cancer-facts-and-figures-2019.pdf>. Accessed February 2019. Ljungberg B, Campbell S and Cho H. The Epidemiology of Renal Cell Carcinoma. *Eur Urol*. 2011;60:615-621. Dolan DE, Gupta S. PD-1 pathway inhibitors: changing the landscape of cancer immunotherapy. *Cancer Control*. 2014;21(3):231-237. Dahan R, Sega E, Engelhardt J, Selby M, Korman AJ, Ravetch JV. FcγRs modulate the anti-tumor activity of antibodies targeting the PD-1/PD-L1 axis. *Cancer Cell*. 2015;28(3):285-295. Boyerinas B, Jochems C, Fantini M, et al. Antibody-dependent cellular cytotoxicity activity of a novel anti-PD-L1 antibody avelumab (MSB0010718C) on human tumor cells. *Cancer Immunol Res*. 2015;3(10):1148-1157. Kohrt HE, Houot R, Marabelle A, et al. Combination strategies to enhance antitumor ADCC. *Immunotherapy*. 2012;4(5):511-527. Hamilton G, Rath B. Avelumab: combining immune checkpoint inhibition and antibody-dependent cytotoxicity. *Expert Opin Biol Ther*. 2017;17(4):515-523.

Merck Media Relations: +1 781 427 4351 [noelle.piscitelli@emdserono.com](mailto:noelle.piscitelli@emdserono.com) Investor Relations: +49 6151 72 3321 [investor.relations@merckgroup.com](mailto:investor.relations@merckgroup.com) Pfizer Media Relations: +1 212 733 6213 [jessica.m.smith@pfizer.com](mailto:jessica.m.smith@pfizer.com) Investor Relations: +1 212 733

8160 ryan.crowe@pfizer.com