

European Commission Approves BAVENCIO® (avelumab) Plus Axitinib Combination for First-Line Treatment of Patients With Advanced Renal Cell Carcinoma

Monday, October 28, 2019 - 04:00pm

Not intended for UK-based media

- **EU approval based on JAVELIN Renal 101 trial results demonstrating significant improvement in progression-free survival with BAVENCIO in combination with axitinib compared with sunitinib**
- **Combination regimen approved across all IMDC prognostic risk groups and irrespective of PD-L1 expression**

Darmstadt, Germany and New York, US, October 28, 2019 – Merck KGaA, Darmstadt, Germany, which operates its biopharmaceutical business as EMD Serono in the US and Canada, and Pfizer Inc. (NYSE: PFE) today announced that the European Commission (EC) has approved BAVENCIO® (avelumab) in combination with axitinib for the first-line treatment of adult patients with advanced renal cell carcinoma (RCC). The approval was based on positive interim results from the Phase III JAVELIN Renal 101 study, which demonstrated that BAVENCIO in combination with axitinib significantly lowered risk of disease progression or death by 31% (HR: 0.69 [95% CI: 0.574–0.825; p<0.0001]) and nearly doubled objective response rate (ORR; 52.5% [95% CI: 47.7-57.2] vs. 27.3% [95% CI: 23.2-31.6]) compared with sunitinib in patients with advanced RCC regardless of PD-L1 status. The study included patients across International Metastatic Renal Cell Carcinoma Database Consortium (IMDC) prognostic risk groups. Improvement in progression-free survival (PFS) was observed across pre-specified subgroups in patients receiving the treatment combination.¹ Merck KGaA, Darmstadt, Germany, and Pfizer have a global strategic alliance to jointly develop and commercialize BAVENCIO.

"There is a high incidence of kidney cancer in Europe, and for the most common type, renal cell carcinoma, we continue to need additional treatment options, particularly for patients with advanced disease, where outcomes are poorest," said Professor James Larkin, Consultant Medical Oncologist at The Royal Marsden NHS Foundation Trust and Professor at the Institute of Cancer Research (ICR). "We've seen a demonstrated efficacy benefit and safety and tolerability profile for avelumab in combination with axitinib across all prognostic risk groups in patients with advanced renal cell carcinoma, so today's approval in Europe brings an important option that can help healthcare professionals optimize treatment strategies across risk stratification."

In 2018, an estimated 136,500 new cases of kidney cancer were diagnosed in Europe, and approximately 54,700 people died from the disease.² Many patients living with advanced RCC do not go on to receive additional treatment after first-line therapy,^{3,4} for reasons that may include poor performance status or adverse events from their initial treatment.^{3,5,6} The five-year survival rate for patients with advanced RCC is approximately 12%.⁷

"This first European approval of an anti-PD-L1 as part of a combination treatment for advanced renal cell carcinoma builds on our commitment to bringing innovative treatment options to patients with hard-to-treat cancers through our extensive JAVELIN clinical trial program," said Rehan Verjee, Global Head of Innovative Medicine Franchises for the Biopharma business of Merck KGaA, Darmstadt, Germany. "RCC is the most common form of kidney cancer, accounting for 90% of diagnoses. We are now working to make BAVENCIO in combination with axitinib available for patients with advanced renal cell carcinoma as quickly as possible."

"The European Commission approval of BAVENCIO in combination with axitinib has the potential to bring even more patients with advanced renal cell carcinoma a new first-line treatment, and it allows us to continue to deliver on our more than decade-long passion to do more for patients with kidney cancer," said Andy Schmeltz, Global President, Pfizer Oncology. "We thank all of the researchers, doctors, advocates, patients and their families who helped get us here today, and we will continue in our fight against this advanced cancer."

The EC's decision follows the U.S. Food and Drug Administration (FDA) approval of BAVENCIO in combination with axitinib for the first-line treatment of patients with advanced RCC in May 2019. A supplemental application for BAVENCIO in combination with axitinib in unresectable or metastatic RCC was submitted in Japan in January 2019.

Additionally, with this approval, the posology section of the Summary of Product Characteristics for BAVENCIO has been updated. The recommended dose of BAVENCIO as monotherapy is 800 mg administered intravenously over 60 minutes every 2 weeks. Administration of BAVENCIO should continue according to the recommended schedule until disease progression or unacceptable toxicity. The recommended dose of BAVENCIO in combination with axitinib is 800 mg administered intravenously over 60 minutes every 2 weeks and axitinib 5 mg orally taken twice daily (12 hours apart) with or without food until disease progression or unacceptable toxicity.¹

Data from JAVELIN Renal 101 Study Supporting Approval

This approval was based on interim data from the Phase III JAVELIN Renal 101 study, a randomized, multicenter, open-label study of BAVENCIO in combination with axitinib in 886 patients with untreated advanced or metastatic RCC with a clear cell component. The study included patients across risk groups (International Metastatic Renal Cell Carcinoma Database Consortium [IMDC]: 21% favorable, 62% intermediate and 16% poor; Memorial Sloan Kettering Cancer Center [MSKCC]: 22% favorable, 65% intermediate and 11% poor). The primary efficacy endpoints were progression-free survival (PFS) as assessed by a Blinded Independent Central Review (BICR) using RECIST v1.1 and overall survival (OS) in the first-line treatment of patients with advanced RCC who have PD-L1-positive tumors (PD L1 expression level ≥1%). PFS based on BICR assessment per RECIST v1.1 and OS irrespective of PD L1 expression, objective response (OR), time to response (TTR), duration of response (DOR) and safety are included as secondary endpoints. The study is continuing for OS.

In the analysis, BAVENCIO in combination with axitinib significantly improved median PFS compared with sunitinib by more than five months in patients irrespective of PD-L1 expression (13.3 months [95% CI: 11.1–15.3] vs. 8.0 months [95% CI: 6.7–9.8]). With a median follow-up for OS of 19 months, data for the trial's other endpoint of OS were immature, with 27% of deaths, and the trial is continuing as planned. The hazard ratio for OS in patients treated with BAVENCIO in combination with axitinib compared with sunitinib was 0.80 (95% CI: 0.616, 1.027) at the interim analysis.

The most common adverse reactions were diarrhea (62.8%), hypertension (49.3%), fatigue (42.9%), nausea (33.5%), dysphonia (32.7%), decreased appetite (26.0%), hypothyroidism (25.2%), cough (23.7%), headache (21.3%), dyspnea (20.9%), and arthralgia (20.9%).

About the JAVELIN Clinical Development Program

The clinical development program for avelumab, known as JAVELIN, involves at least 30 clinical programs and more than 10,000 patients evaluated across more than 15 different tumor types. In addition to RCC, these tumor types include gastric/gastro-esophageal junction cancer, head and neck cancer, Merkel cell carcinoma, non-small cell lung cancer and urothelial carcinoma.

About BAVENCIO® (avelumab)

BAVENCIO is a human anti-programmed death ligand-1 (PD-L1) antibody. BAVENCIO has been shown in preclinical models to engage both the adaptive and innate immune functions. By blocking the interaction of PD-L1 with PD-1 receptors, BAVENCIO has been shown to release the suppression of the T cell-mediated antitumor immune response in preclinical models.⁸⁻¹⁰ In November 2014, Merck KGaA, Darmstadt, Germany, and Pfizer announced a strategic alliance to co-develop and co-commercialize BAVENCIO.

BAVENCIO Approved Indications in the US

BAVENCIO® (avelumab) in combination with axitinib is indicated in the US for the first-line treatment of patients with advanced renal cell carcinoma (RCC).

In the US, the FDA granted accelerated approval for BAVENCIO for the treatment of (i) adults and pediatric patients 12 years and older with metastatic Merkel cell carcinoma (mMCC) and (ii) patients with locally advanced or metastatic urothelial carcinoma (mUC) who have disease progression during or following platinum-containing chemotherapy, or have disease progression within 12 months of neoadjuvant or adjuvant treatment with platinum-containing chemotherapy. These indications are approved under accelerated approval based on tumor response rate and duration of response. Continued approval for these indications may be contingent upon verification and description of clinical benefit in confirmatory trials.

Avelumab is currently approved for patients with MCC in 50 countries globally, with the majority of these approvals in a broad indication that is not limited to a specific line of treatment.

BAVENCIO Important Safety Information from the US FDA-Approved Label

BAVENCIO can cause **immune-mediated pneumonitis**, including fatal cases. Monitor patients for signs and symptoms of pneumonitis, and evaluate suspected cases with radiographic imaging. Administer corticosteroids for Grade 2 or greater pneumonitis. Withhold BAVENCIO for moderate (Grade 2) and permanently discontinue for severe (Grade 3), life-threatening (Grade 4), or recurrent moderate (Grade 2) pneumonitis. Pneumonitis occurred in 1.2% of patients, including one (0.1%) patient with Grade 5, one (0.1%) with Grade 4, and five (0.3%) with Grade 3.

BAVENCIO can cause **hepatotoxicity and immune-mediated hepatitis**, including fatal cases. Monitor patients for abnormal liver tests prior to and periodically during treatment. Administer corticosteroids for Grade 2 or greater hepatitis. Withhold BAVENCIO for moderate (Grade 2) immune-mediated hepatitis until resolution and permanently discontinue for severe (Grade 3) or life-threatening (Grade 4) immune-mediated hepatitis. Immune-mediated hepatitis occurred with BAVENCIO as a single agent in 0.9% of patients, including two (0.1%) patients with Grade 5, and 11 (0.6%) with Grade 3.

BAVENCIO in combination with axitinib can cause **hepatotoxicity** with higher than expected frequencies of Grade 3 and 4 alanine aminotransferase (ALT) and aspartate aminotransferase (AST) elevation. Consider more

frequent monitoring of liver enzymes as compared to when the drugs are used as monotherapy. Withhold BAVENCIO and axitinib for moderate (Grade 2) hepatotoxicity and permanently discontinue the combination for severe or life-threatening (Grade 3 or 4) hepatotoxicity. Administer corticosteroids as needed. In patients treated with BAVENCIO in combination with axitinib, Grades 3 and 4 increased ALT and AST occurred in 9% and 7% of patients, respectively, and immune-mediated hepatitis occurred in 7% of patients, including 4.9% with Grade 3 or 4.

BAVENCIO can cause **immune-mediated colitis**. Monitor patients for signs and symptoms of colitis. Administer corticosteroids for Grade 2 or greater colitis. Withhold BAVENCIO until resolution for moderate or severe (Grade 2 or 3) colitis until resolution. Permanently discontinue for life-threatening (Grade 4) or recurrent (Grade 3) colitis upon reinitiation of BAVENCIO. Immune-mediated colitis occurred in 1.5% of patients, including seven (0.4%) with Grade 3.

BAVENCIO can cause **immune-mediated endocrinopathies**, including adrenal insufficiency, thyroid disorders, and type 1 diabetes mellitus.

Monitor patients for signs and symptoms of **adrenal insufficiency** during and after treatment, and administer corticosteroids as appropriate. Withhold BAVENCIO for severe (Grade 3) or life-threatening (Grade 4) adrenal insufficiency. Adrenal insufficiency was reported in 0.5% of patients, including one (0.1%) with Grade 3.

Thyroid disorders can occur at any time during treatment. Monitor patients for changes in thyroid function at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation. Manage hypothyroidism with hormone replacement therapy and hyperthyroidism with medical management. Withhold BAVENCIO for severe (Grade 3) or life-threatening (Grade 4) thyroid disorders. Thyroid disorders, including hypothyroidism, hyperthyroidism, and thyroiditis, were reported in 6% of patients, including three (0.2%) with Grade 3.

Type 1 diabetes mellitus including diabetic ketoacidosis: Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Withhold BAVENCIO and administer antihyperglycemics or insulin in patients with severe or life-threatening (Grade 3) hyperglycemia, and resume treatment when metabolic control is achieved. Type 1 diabetes mellitus without an alternative etiology occurred in 0.1% of patients, including two cases of Grade 3 hyperglycemia.

BAVENCIO can cause **immune-mediated nephritis and renal dysfunction**. Monitor patients for elevated serum creatinine prior to and periodically during treatment. Administer corticosteroids for Grade 2 or greater nephritis. Withhold BAVENCIO for moderate (Grade 2) or severe (Grade 3) nephritis until resolution to Grade 1 or lower. Permanently discontinue BAVENCIO for life-threatening (Grade 4) nephritis. Immune-mediated nephritis occurred in 0.1% of patients.

BAVENCIO can result in **other severe and fatal immune-mediated adverse reactions** involving any organ system during treatment or after treatment discontinuation. For suspected immune-mediated adverse reactions, evaluate to confirm or rule out an immune-mediated adverse reaction and to exclude other causes. Depending on the severity of the adverse reaction, withhold or permanently discontinue BAVENCIO, administer high-dose corticosteroids, and initiate hormone replacement therapy, if appropriate. Resume BAVENCIO when the immune-mediated adverse reaction remains at Grade 1 or lower following a corticosteroid taper. Permanently discontinue BAVENCIO for any severe (Grade 3) immune-mediated adverse reaction that recurs and for any life-threatening (Grade 4) immune-mediated adverse reaction. The following clinically significant immune-mediated adverse reactions occurred in less than 1% of 1738 patients treated with BAVENCIO as a single agent or in 489 patients who received BAVENCIO in combination with axitinib: myocarditis including fatal cases, pancreatitis including fatal cases, myositis, psoriasis, arthritis, exfoliative dermatitis, erythema multiforme, pemphigoid, hypopituitarism, uveitis, Guillain-Barré syndrome, and systemic inflammatory response.

BAVENCIO can cause severe or life-threatening **infusion-related reactions**. Premedicate patients with an antihistamine and acetaminophen prior to the first 4 infusions and for subsequent infusions based upon clinical judgment and presence/severity of prior infusion reactions. Monitor patients for signs and symptoms of infusion-related reactions, including pyrexia, chills, flushing, hypotension, dyspnea, wheezing, back pain, abdominal pain, and urticaria. Interrupt or slow the rate of infusion for mild (Grade 1) or moderate (Grade 2) infusion-related reactions. Permanently discontinue BAVENCIO for severe (Grade 3) or life-threatening (Grade 4) infusion-related reactions. Infusion-related reactions occurred in 25% of patients, including three (0.2%) patients with Grade 4 and nine (0.5%) with Grade 3.

BAVENCIO in combination with axitinib can cause **major adverse cardiovascular events (MACE)** including severe and fatal events. Consider baseline and periodic evaluations of left ventricular ejection fraction. Monitor for signs and symptoms of cardiovascular events. Optimize management of cardiovascular risk factors, such as hypertension, diabetes, or dyslipidemia. Discontinue BAVENCIO and axitinib for Grade 3-4 cardiovascular events. MACE occurred in 7% of patients with advanced RCC treated with BAVENCIO in combination with axitinib compared to 3.4% treated with sunitinib. These events included death due to cardiac events (1.4%), Grade 3-4 myocardial infarction (2.8%), and Grade 3-4 congestive heart failure (1.8%).

BAVENCIO can cause **fetal harm** when administered to a pregnant woman. Advise patients of the potential risk to a fetus including the risk of fetal death. Advise females of childbearing potential to use effective contraception during treatment with BAVENCIO and for at least 1 month after the last dose of BAVENCIO. It is not known whether BAVENCIO is excreted in human milk. Advise a lactating woman **not to breastfeed** during treatment and for at least 1 month after the last dose of BAVENCIO due to the potential for serious adverse reactions in breastfed infants.

The most common adverse reactions (all grades, ? 20%) in patients with metastatic Merkel cell carcinoma (MCC) were fatigue (50%), musculoskeletal pain (32%), diarrhea (23%), nausea (22%), infusion-related reaction (22%), rash (22%), decreased appetite (20%), and peripheral edema (20%).

Selected treatment-emergent laboratory abnormalities (all grades, ? 20%) in patients with metastatic MCC were lymphopenia (49%), anemia (35%), increased aspartate aminotransferase (34%), thrombocytopenia (27%), and increased alanine aminotransferase (20%).

The most common adverse reactions (all grades, ? 20%) in patients with locally advanced or metastatic urothelial carcinoma (UC) were fatigue (41%), infusion-related reaction (30%), musculoskeletal pain (25%), nausea (24%), decreased appetite/hypophagia (21%), and urinary tract infection (21%).

Selected laboratory abnormalities (Grades 3-4, ? 3%) in patients with locally advanced or metastatic UC were hyponatremia (16%), increased gamma-glutamyltransferase (12%), lymphopenia (11%), hyperglycemia (9%), increased alkaline phosphatase (7%), anemia (6%), increased lipase (6%), hyperkalemia (3%), and increased aspartate aminotransferase (3%).

Fatal adverse reactions occurred in 1.8% of patients with advanced renal cell carcinoma (RCC) receiving BAVENCIO in combination with axitinib. These included sudden cardiac death (1.2%), stroke (0.2%), myocarditis (0.2%), and necrotizing pancreatitis (0.2%).

The most common adverse reactions (all grades, ?20%) in patients with advanced RCC receiving BAVENCIO in combination with axitinib (vs sunitinib) were diarrhea (62% vs 48%), fatigue (53% vs 54%), hypertension (50% vs 36%), musculoskeletal pain (40% vs 33%), nausea (34% vs 39%), mucositis (34% vs 35%), palmar-plantar erythrodysesthesia (33% vs 34%), dysphonia (31% vs 3.2%), decreased appetite (26% vs 29%), hypothyroidism (25% vs 14%), rash (25% vs 16%), hepatotoxicity (24% vs 18%), cough (23% vs 19%), dyspnea (23% vs 16%), abdominal pain (22% vs 19%), and headache (21% vs 16%).

Selected laboratory abnormalities (all grades, 72%) worsening from baseline in patients with advanced RCC receiving BAVENCIO in combination with axitinib (vs sunitinib) were blood triglycerides increased (71% vs 48%), blood creatinine increased (62% vs 68%), blood cholesterol increased (57% vs 22%), alanine aminotransferase increased (ALT) (50% vs 46%), aspartate aminotransferase increased (AST) (47% vs 57%), blood sodium decreased (38% vs 37%), lipase increased (37% vs 25%), blood potassium increased (35% vs 28%), platelet count decreased (27% vs 80%), blood bilirubin increased (21% vs 23%), and hemoglobin decreased (21% vs 65%).

Please see full [US Prescribing Information](#) and [Medication Guide](#) available at <http://www.BAVENCIO.com>."

Axitinib Important Safety Information from the US FDA-Approved Label

Hypertension including **hypertensive crisis** has been observed with axitinib. Blood pressure should be well controlled prior to initiating axitinib. Monitor for hypertension and treat as needed. For persistent hypertension, despite use of antihypertensive medications, reduce the dose. Discontinue axitinib if hypertension is severe and persistent despite use of antihypertensive therapy and dose reduction of axitinib, and discontinuation should be considered if there is evidence of hypertensive crisis.

Arterial and venous thrombotic events have been observed with axitinib and can be fatal. Use with caution in patients who are at increased risk or who have a history of these events.

Hemorrhagic events, including fatal events, have been reported with axitinib. Axitinib has not been studied in patients with evidence of untreated brain metastasis or recent active gastrointestinal bleeding and should not be used in those patients. If any bleeding requires medical intervention, temporarily interrupt the axitinib dose.

Cardiac failure has been observed with axitinib and can be fatal. Monitor for signs or symptoms of cardiac failure throughout treatment with axitinib. Management of cardiac failure may require permanent discontinuation of axitinib.

Gastrointestinal perforation and fistula, including death, have occurred with axitinib. Use with caution in patients at risk for gastrointestinal perforation or fistula. Monitor for symptoms of gastrointestinal perforation or fistula periodically throughout treatment.

Hypothyroidism requiring thyroid hormone replacement has been reported with axitinib. Monitor thyroid function before initiation of, and periodically throughout, treatment.

No formal studies of the effect of axitinib on **wound healing** have been conducted. Stop axitinib at least 24 hours prior to scheduled surgery.

Reversible Posterior Leukoencephalopathy Syndrome (RPLS) has been observed with axitinib. If signs or symptoms occur, permanently discontinue treatment.

Proteinuria has been observed with axitinib. Monitor for proteinuria before initiation of, and periodically throughout, treatment with axitinib. For moderate to severe proteinuria, reduce the dose or temporarily interrupt treatment.

Liver enzyme elevation has been observed during treatment with axitinib. Monitor ALT, AST, and bilirubin before initiation of, and periodically throughout, treatment.

For patients with moderate **hepatic impairment**, the starting dose should be decreased. Axitinib has not been studied in patients with severe hepatic impairment.

Axitinib can cause **fetal harm**. Advise patients of the potential risk to the fetus and to use effective contraception during treatment.

Avoid strong **CYP3A4/5 inhibitors**. If unavoidable, reduce the dose. Grapefruit or grapefruit juice may also increase axitinib plasma concentrations and should be avoided.

Avoid strong **CYP3A4/5 inducers** and, if possible, avoid moderate CYP3A4/5 inducers.

Please see full [Prescribing Information](#) for axitinib.

ADVERSE REACTIONS (BAVENCIO + AXITINIB)

Fatal adverse reactions occurred in 1.8% of patients with **advanced renal cell carcinoma (RCC)** receiving BAVENCIO in combination with axitinib. These included sudden cardiac death (1.2%), stroke (0.2%), myocarditis (0.2%), and necrotizing pancreatitis (0.2%).

The most common adverse reactions (all grades, ≥20%) in patients with advanced RCC receiving BAVENCIO in combination with axitinib (vs sunitinib) were diarrhea (62% vs 48%), fatigue (53% vs 54%), hypertension (50% vs 36%), musculoskeletal pain (40% vs 33%), nausea (34% vs 39%), mucositis (34% vs 35%), palmar-plantar erythrodysesthesia (33% vs 34%), dysphonia (31% vs 3.2%), decreased appetite (26% vs 29%), hypothyroidism (25% vs 14%), rash (25% vs 16%), hepatotoxicity (24% vs 18%), cough (23% vs 19%), dyspnea (23% vs 16%), abdominal pain (22% vs 19%), and headache (21% vs 16%).

Selected laboratory abnormalities (all grades, ≥20%) worsening from baseline in patients with **advanced RCC** receiving BAVENCIO in combination with axitinib (vs sunitinib) were blood triglycerides increased (71% vs 48%), blood creatinine increased (62% vs 68%), blood cholesterol increased (57% vs 22%), alanine aminotransferase increased (ALT) (50% vs 46%), aspartate aminotransferase increased (AST) (47% vs 57%), blood sodium decreased (38% vs 37%), lipase increased (37% vs 25%), blood potassium increased (35% vs 28%), platelet count decreased (27% vs 80%), blood bilirubin increased (21% vs 23%), and hemoglobin decreased (21% vs 65%).

The most common adverse reactions (all grades, ≥20%) in patients with **advanced RCC** receiving BAVENCIO in combination with axitinib (vs sunitinib) were diarrhea (62% vs 48%), fatigue (53% vs 54%), hypertension (50% vs 36%), musculoskeletal pain (40% vs 33%), nausea (34% vs 39%), mucositis (34% vs 35%), palmar-plantar erythrodysesthesia (33% vs 34%), dysphonia (31% vs 3.2%), decreased appetite (26% vs 29%), hypothyroidism (25% vs 14%), rash (25% vs 16%), hepatotoxicity (24% vs 18%), cough (23% vs 19%), dyspnea (23% vs 16%), abdominal pain (22% vs 19%), and headache (21% vs 16%).

Selected laboratory abnormalities (all grades, ≥20%) worsening from baseline in patients with **advanced RCC** receiving BAVENCIO in combination with axitinib (vs sunitinib) were blood triglycerides increased (71% vs 48%), blood creatinine increased (62% vs 68%), blood cholesterol increased (57% vs 22%), alanine aminotransferase increased (ALT) (50% vs 46%), aspartate aminotransferase increased (AST) (47% vs 57%), blood sodium decreased (38% vs 37%), lipase increased (37% vs 25%), blood potassium increased (35% vs 28%), platelet count decreased (27% vs 80%), blood bilirubin increased (21% vs 23%), and hemoglobin decreased (21% vs 65%).

About Merck KGaA, Darmstadt, Germany-Pfizer Alliance

Immuno-oncology is a top priority for Merck KGaA, Darmstadt, Germany and Pfizer. The global strategic alliance between Merck KGaA, Darmstadt, Germany and Pfizer enables the companies to benefit from each other's strengths and capabilities and further explore the therapeutic potential of BAVENCIO, an anti-PD-L1 antibody initially discovered and developed by Merck KGaA, Darmstadt, Germany. The immuno-oncology alliance is jointly developing and commercializing BAVENCIO. The alliance is focused on developing high-priority international clinical programs to investigate BAVENCIO as a monotherapy as well as combination regimens, and is striving to find new ways to treat cancer.

All Merck KGaA, Darmstadt, Germany, press releases are distributed by e-mail at the same time they become available on the EMD Group Website. In case you are a resident of the USA or Canada please go to www.emdgroup.com/subscribe to register again for your online subscription of this service as our newly introduced geo-targeting requires new links in the email. You may later change your selection or discontinue this service.

About Merck KGaA, Darmstadt, Germany

Merck KGaA, Darmstadt, Germany, a leading science and technology company, operates across healthcare, life science and performance materials. Around 56,000 employees work to make a positive difference to millions of people's lives every day by creating more joyful and sustainable ways to live. From advancing gene editing technologies and discovering unique ways to treat the most challenging diseases to enabling the intelligence of devices – the company is everywhere. In 2018, Merck KGaA, Darmstadt, Germany, generated sales of € 14.8 billion in 66 countries.

The company holds the global rights to the name and trademark "Merck" internationally. The only exceptions are the United States and Canada, where the business sectors of Merck KGaA, Darmstadt, Germany operate as EMD Serono in healthcare, MilliporeSigma in life science, and EMD Performance Materials. Since its founding 1668, scientific exploration and responsible entrepreneurship have been key to the company's technological and scientific advances. To this day, the founding family remains the majority owner of the publicly listed company.

Pfizer Inc.: Breakthroughs that change patients' lives

At Pfizer, we apply science and our global resources to bring therapies to people that extend and significantly improve their lives. We strive to set the standard for quality, safety and value in the discovery, development and manufacture of health care products, including innovative medicines and vaccines. Every day, Pfizer colleagues work across developed and emerging markets to advance wellness, prevention, treatments and cures that challenge the most feared diseases of our time. Consistent with our responsibility as one of the world's premier innovative biopharmaceutical companies, we collaborate with health care providers, governments and local communities to support and expand access to reliable, affordable health care around the world. For more than 150 years, we have worked to make a difference for all who rely on us. We routinely post information that may be important to investors on our website at www.pfizer.com. In addition, to learn more, please visit us on www.pfizer.com and follow us on Twitter at [@Pfizer](https://twitter.com/Pfizer) and [@Pfizer_News](https://twitter.com/Pfizer_News), [LinkedIn](https://www.linkedin.com/company/pfizer), [YouTube](https://www.youtube.com/pfizer) and like us on Facebook at [Facebook.com/Pfizer](https://www.facebook.com/Pfizer).

Pfizer Disclosure Notice

The information contained in this release is as of October 28, 2019. Pfizer assumes no obligation to update forward-looking statements contained in this release as the result of new information or future events or developments.

This release contains forward-looking information about BAVENCIO (avelumab), including a new indication approved in the European Union for BAVENCIO in combination with axitinib for the treatment of patients with advanced renal cell carcinoma, the alliance between Merck KGaA, Darmstadt, Germany, and Pfizer involving BAVENCIO and clinical development plans, including their potential benefits, that involves substantial risks and uncertainties that could cause actual results to differ materially from those expressed or implied by such statements. Risks and uncertainties include, among other things, uncertainties regarding the commercial success of BAVENCIO and axitinib; the uncertainties inherent in research and development, including the ability to meet anticipated clinical endpoints, commencement and/or completion dates for our clinical trials, regulatory submission dates, regulatory approval dates and/or launch dates, as well as the possibility of unfavorable new clinical data and further analyses of existing clinical data; risks associated with interim data; the risk that clinical trial data are subject to differing interpretations and assessments by regulatory authorities; whether regulatory authorities will be satisfied with the design of and results from our clinical studies; whether and when any drug applications may be filed for BAVENCIO in combination with axitinib in any other jurisdictions or in any jurisdictions for any other potential indications for BAVENCIO or combination therapies; whether and when the pending application in Japan for BAVENCIO in combination with axitinib may be approved and whether and when regulatory authorities in any jurisdictions where any other applications are pending or may be submitted for BAVENCIO or combination therapies, including BAVENCIO in combination with axitinib may approve any such applications, which will depend on myriad factors, including making a determination as to whether the product's benefits outweigh its known risks and determination of the product's efficacy, and, if approved, whether they will be commercially successful; decisions by regulatory authorities impacting labeling, manufacturing processes, safety and/or other matters that could affect the availability or commercial potential of BAVENCIO or combination therapies, including BAVENCIO in combination with axitinib; and competitive developments.

A further description of risks and uncertainties can be found in Pfizer's Annual Report on Form 10-K for the fiscal year ended December 31, 2018, and in its subsequent reports on Form 10-Q, including in the sections thereof captioned "Risk Factors" and "Forward-Looking Information and Factors That May Affect Future Results", as well as in its subsequent reports on Form 8-K, all of which are filed with the U.S. Securities and Exchange Commission and available at www.sec.gov and www.pfizer.com.

References

1. BAVENCIO® (avelumab) EU SmPC. Available from: <http://www.ema.europa.eu/ema/>. Accessed October 2019.
2. Ferlay J, Colombet M, Soerjomataram I, et al. Cancer incidence and mortality patterns in Europe: Estimates for 40 countries and 25 major cancers in 2018. *Eur J Cancer*. 2018;103:356-387.
3. Eggers H, Ivanyi P, Hornig M, Grünwald V. Predictive factors for second-line therapy in metastatic renal cell carcinoma: a retrospective analysis. *J Kidney Cancer VHL*. 2017;4(1):8-15.
4. Motzer R, et al. Nivolumab plus Ipilimumab versus Sunitinib in Advanced Renal-Cell Carcinoma. *The New England Journal of Medicine*. 2018;378:1277-1290.
5. Eichelberg C, Vervenne WL, De Santis M, et al. SWITCH: A randomised, sequential, open-label study to evaluate the efficacy and safety of sorafenib-sunitinib versus sunitinib-sorafenib in the treatment of metastatic renal cell cancer. *Eur Urol*. 2015;68:837-847.
6. Motzer RJ, Barrios CH, Kim TM, et al. Phase II randomized trial comparing sequential first-line everolimus and second-line sunitinib versus first-line sunitinib and second-line everolimus in patients with metastatic renal cell carcinoma. *J Clin Oncol*. 2014;32:2765-2772.
7. Ridge C, Pua B, Madoff D. Epidemiology and staging of renal cell carcinoma. *Semin Intervent Radiol*. 2014;31(1):3-8.

8. Dolan DE, Gupta S. PD-1 pathway inhibitors: changing the landscape of cancer immunotherapy. *Cancer Control*. 2014;21(3):231-237.
9. Dahan R, Segal E, Engelhardt J, et al. FcγRs modulate the anti-tumor activity of antibodies targeting the PD-1/PD-L1 axis. *Cancer Cell*. 2015;28(3):285-295.
10. Boyerinas B, Jochems C, Fantini M, et al. Antibody-dependent cellular cytotoxicity activity of a novel anti-PD-L1 antibody avelumab (MSB0010718C) on human tumor cells. *Cancer Immunol Res*. 2015;3(10):1148-1157.

Your Contacts Merck KGaA, Darmstadt, Germany Media Relations +49 6151 72 6328
friederike.segeberg@emdgroup.com Investor Relations +49 6151 72-3321
investor.relations@emdgroup.com Pfizer Inc., New York, USA Media Jessica Smith Lisa O'Neill +1 212 733
6213 +44 7929 339 560 Investor Relations Ryan Crowe +1 212 733 8160